

## SCIENTIFIC NOTE

**NEOALLOCERAEA NOM. NOV., A REPLACEMENT NAME  
FOR THE SUBGENUS *ALLOCERAEA* BENICK, 1934  
(COLEOPTERA: STAPHYLINIDAE)  
NON SCHULZE, 1918<sup>1</sup>**

Hüseyin Özdkmen<sup>2</sup> and Mustafa C. Darılmaz<sup>2</sup>

In an effort to reduce the number of homonyms in Staphylinidae, we found a genus group name which had been previously published for another taxon, making it a junior homonym. In accordance with Article 60 of the International Code of Zoological Nomenclature (1999), we propose a substitute name for this genus name.

**Family Staphylinidae**

**Genus *Plataraea* Thomson, 1858**

**Subgenus *Neoalloceraea* nom. nov.**

*Alloceraea* Benick, 1934. Ent. Blätter, 31, 164. (Coleoptera: Staphylinidae). Preoccupied by *Alloceraea* Schulze, 1918. S.B. Ges. naturf. Freunde Berlin, 1918, 62 (Acari: Ixodida).

Benick (1934) proposed the subgenus *Alloceraea* (type species, *Atheta fiorii* Bernhauer, 1908 by original designation) and placed it in the rove beetle genus *Plataraea* Thomson, 1858 (Coleoptera: Staphyliniformia: Staphylinoidea: Staphylinidae: Aleocharinae: Athetini: *Plataraea*). *Alloceraea* is still used as a valid genus group in the Coleoptera (e.g. Löbl and Smetana, 2004). Currently *Alloceraea* includes only one species, *Alloceraea fiorii* (Bernhauser, 1907) (formerly *Atheta fiorii* Bernhauser, 1908) as the type, from Italy.

Unfortunately, the generic name, *Alloceraea*, was already preoccupied by Schulze (1918), who had described *Alloceraea* (type species *Haemaphysalis inermis* Birula, 1895) as a genus of tick (Acari: Ixodida: Ixodina: Amblyommidae: Rhipicephalinae). Currently, *Alloceraea* is considered a subgenus of the speciose genus *Haemaphysalis* Koch, 1844.

Thus, the subgenus name *Alloceraea* Benick, 1934 is a junior homonym of the subgenus name *Alloceraea* Schulze, 1918. Under the Zoological Code (ICZN, 1999), it must be rejected and replaced, and so a new replacement name *Neoalloceraea* nom. nov. for *Alloceraea* Benick, 1934 is herein proposed in the present work.

<sup>1</sup> Submitted on September 17, 2007. Accepted on October 18, 2007.

<sup>2</sup> Gazi University, Arts and Sciences Faculty, Department of Biology, 06500-Teknikokullar Ankara, Turkey. E-mails: ozdikmen@gazi.edu.tr, mdarilmaz@yahoo.com (corresponding author), respectively.

Etymology: From the preexisting subgenus name *Alloceraea*.

Distribution: Europe (Italy).

**Summary of nomenclatural changes:**

**Genus** *Plataraea* Thomson, 1858

**Subgenus** *Plataraea (Neoalloceraea)* nom. nov.

= *Plataraea (Alloceraea)* Benick, 1934 (non Schulze, 1918)

Species *Plataraea (Neoalloceraea) fiorii* (Bernhauer, 1908) **comb. nov.**

= *Atheta fiorii* Bernhauer, 1908

= *Plataraea (Alloceraea) fiorii* (Bernhauer, 1908)

**LITERATURE CITED**

- Benick, G.** 1934. Revision der Untergattungen *Plataraea* Thoms. Und *Aerostiba* Bernh. (Gattung *Atheta*, Staphyl.) Entomologische Blätter 31: 161-166.
- ICZN.** 1999. International Code of Zoological Nomenclature. Fourth Edition. The International Trust for Zoological Nomenclature, London.
- Löbl, I. and A. Smetana.** 2004. Catalogue of Palaearctic Coleoptera, Vol. 2. Apollo Books 942 pp.
- Schulze, P.** 1918. Ein Beitrag zur Zeckenfauna Mazedoniens. Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin 1918 (2): 61-66.